

Ro52 ANTIGEN

Product Datasheet

Description of the Product:

Recombinant full-length human sequence protein. After coating onto ELISA plates the product will bind autoantibodies to Ro52.

Purity: The Ro52 autoantigen (52 kDa) is more than 90% pure, as assessed by SDS-polyacrylamide gel electrophoresis.

Storage & Handling:

Store at -65°C or below (long term). Avoid repeated freezing and thawing. CENP-B buffer contains SDS, this must be completely solubilized before use:

- Thaw at room temperature for at least an hour, OR
- Warm the vial above 19°C, OR
- Place the vial in a 37°C water bath for 5 minutes.

Gentle but thorough mixing is required before use.

Clinical and Biochemical Data:

Autoantibodies to the Ro(SSA) antigen are one of the most frequent serological autoimmune markers in rheumatic diseases^{1,2} (refer also to AroTec product ATR02 Ro(SSA) antigen). Anti-Ro-positive sera may contain two different types of autoantibody; those directed to a 60 kDa and those directed to a 52 kDa polypeptide component (referred to as Ro60 and Ro52 respectively)³. While the vast majority of Ro(SSA) positive sera react with both of these components, anti-Ro60 antibody has been reported to occur without anti-Ro52 antibody only in SLE sera^{4,5} whereas anti-Ro52 antibody has been reported to occur in the absence of anti-Ro60 antibody in idiopathic inflammatory myopathy⁶, dermatomyositis^{7,8} and scleroderma^{7,8}. Antibodies to Ro52 have been observed to occur not only monospecifically or with Ro(SSA)/La(SSB), but also concurrently with antibodies to Jo-1⁶ and to other aminoacyl transferases⁷, with anti-SRP⁷, anti-PMScl⁷, anti-CENP-B⁹, anti-RNP⁹ and antihistones⁹. The differing serological and clinical associations of anti-Ro52 and anti-Ro60 suggest that anti-Ro52 should be considered as an additional connective tissue disease serum marker⁹, independent of anti-Ro60(SSA).

Ro52 is a member of the tripartite motif (TRIM) family of proteins. The TRIM motif includes three zinc-binding domains, a RING finger, a B-box type 1 and a B-box type 2, and a centre coiled-coil region (leucine zipper). TRIM proteins are believed to identify specific cell compartments through a process of homomultimerisation^{10,11}. Other functional studies have suggested a role for a putative leucine zipper domain of Ro52 in protein dimer formation and inhibition of transcription activity¹², and involvement of Ro52 in the ubiquitin pathway^{13,14}. Although autoantibodies are known to bind to multiple epitopes on the Ro52 protein¹⁵⁻¹⁷, the leucine zipper domain and an area on the N-terminal side of it have been shown to constitute a prominent linear epitope^{19,21}. Antibodies to the leucine zipper domain of Ro52 have been reported to bind to Ro60 in a conformationally-dependent fashion^{22,23} suggesting there might be some degree of epitope spreading between Ro60 and Ro52.

AROTEC's Ro52 antigen has been expressed and purified in a denatured form – denatured Ro52 is known to react well with Ro52-specific autoantibodies^{22,24}. The use of recombinant antigen for the detection of anti-Ro52 antibodies has been described elsewhere^{25,26}.

Ordering information:

ATR05-02 Ro52 antigen 0.20 mg

ATR05-10 Ro52 antigen 1.0 mg

Custom pack sizes available on request

Methodology:

The following is an ELISA procedure which can be used to detect anti-Ro52 autoantibodies in human serum using the ATR05 purified Ro52 autoantigen:

1. Dilute the purified antigen to 0.2-0.5 µg/ml in PBS (10 mM potassium phosphate, pH 7.4, 0.15 M NaCl).
2. Coat ELISA plates with 100 µl of diluted antigen per well. Cover and incubate 24 hours at +4°C.
3. Empty the plates and remove excess liquid by tapping on a paper towel.
4. Block excess protein binding sites by adding 200 µl PBS containing 1% BSA per well. Cover and incubate at +4°C overnight.
5. Empty plates and apply 100 µl of serum samples diluted 1:100 in PBS / 1% BSA / 1% casein / 0.1% Tween® 20. Incubate at room temperature for 1 hour.
6. Empty plates and add 200 µl PBS / 0.1% Tween® 20 per well. Incubate 5 minutes then empty plates. Repeat this step twice.
7. Apply 100 µl anti-human IgG-enzyme conjugate (horseradish peroxidase or alkaline phosphatase) diluted in PBS / 1% BSA / 1% casein / 0.1% Tween® 20 per well and incubate for 1 hour.
8. Repeat step 6.
9. Add enzyme substrate and stop the reaction when appropriate.
10. Read absorbance in an ELISA spectrophotometer.

References:

1. Tan, E.M. (1989) *Adv. Immunol.* **44**, 93
2. McCauliffe, D.P. & Sontheimer, R.D. (1993) *J. Invest. Dermatol.* **100**, 735
3. Chan, E.K.L. & Buyon, J.P. (1994) *Man. Biol. Markers Dis.* (Kluwer Acad. Publ.) **B4**,1/1-18
4. Ben-Cherit, E. et al. (1990) *Arthritis Rheum.* **33**, 349
5. Slobbe, R.L. et al. (1991) *Clin. Exp. Immunol.* **86**, 99
6. Rutjes, S.A. et al. (1997) *Clin. Exp. Immunol.* **109**, 32
7. Frank, M.B. et al. (1999) *J. Autoimmun.* **12**, 137
8. Peene, I. et al. (2002) *Ann. Rheum. Dis.* **61**, 1099
9. Peene, I. et al. (2002) *Ann. Rheum. Dis.* **61**, 929
10. Raymond, A. et al. (2001) *EMBO J.* **20**, 2140
11. Rhodes, D.A. et al. (2002) *Immunol.* **106**, 246
12. Wang, D. et al. (2001) *Biochim. Biophys. Acta* **1568**, 155
13. Di Donato, F. et al. (2001) *Int. J. Biochem. Cell Biol.* **33**, 924
14. Fukuda-Kamitani, T. & Kamitani, T. (2002) *Biochem. Biophys. Res. Comm.* **292**, 774
15. Bozic, B. et al. (1993) *Clin. Exp. Immunol.* **94**, 227
16. McCauliffe, D.P. et al. (1994) *J. Rheumatol.* **21**, 1073
17. Ricchiuti, V. et al. (1994) *Clin. Exp. Immunol.* **95**, 397
18. Frank, M.B. et al. (1994) *Clin. Exp. Immunol.* **95**, 390
19. St Clair, E.V. et al. (1994) *Arthritis Rheum.* **37**, 1373
20. Blange, I. et al. (1994) *J. Autoimmunity* **7**, 263
21. Peek, R. et al. (1994) *J. Immunol.* **153**, 4321
22. Itoh, Y. et al. (1992) *Autoimmunity* **14**, 89
23. Kurien, B.T. et al. (2001) *Scan. J. Immunol.* **53**, 268
24. Ben-Cherit, E. et al. (1988) *J. Exp. Med.* **167**, 1560
25. McCauliffe, D.P. et al. (1997) *J. Rheumatol.* **24**, 860
26. Pourmand, N. et al. (2000) *Scan. J. Rheumatol.* **29**, 116

